

STAR™ total ankle



STAR has the lowest published rates
of progressive subtalar joint arthritis¹

Device	Patients reporting post-op progressive subtalar joint arthritis	Average follow-up
STAR	20% ¹	9.1 years
Salto-Talaris	43% ²	3.0 years
Fusion	91% ³	22.0 years

Proven.

Visit footankle.stryker.com for more information.

References

1. Jeffrey A. Mann, Roger A. Mann, and Eric Horton. STAR Ankle: Long-Term Results . *Foot & Ankle International* 2011 32: 473-484.
2. Chao J, Choi JH, Gear BJ, Tenenbaum S, Bariteau JT, Brodsky JW. Early radiographic and clinical results of the Salto total ankle arthroplasty as a fixed-bearing device. *Foot and Ankle Surgery* 21 (2015) 91-96.
3. Coester LM, Saltzman CL, Leupold J, Pontarelli W. Long-Term Results Following Ankle Arthrodesis for Post-Traumatic Arthritis. *J Bone Joint Surg Am* 2001;83A(2):219-228.

A surgeon must always rely on his or her own professional clinical judgment when deciding whether to use a particular product when treating a particular patient. Stryker does not dispense medical advice and recommends that surgeons be trained in the use of any particular product before using it in surgery. The information presented is intended to demonstrate the breadth of Stryker product offerings. A surgeon must always refer to the package insert, product label and/or instructions for use before using any Stryker product. Products may not be available in all markets because product availability is subject to the regulatory and/or medical practices in individual markets. Please contact your Stryker representative if you have questions about the availability of Stryker products in your area. Stryker Corporation or its divisions or other corporate affiliated entities own, use or have applied for the following trademarks or service marks: STAR, Stryker. All other trademarks are trademarks of their respective owners or holders.