

# Mako<sup>®</sup> Partial Knee arthroplasty: clinical summary



## Mako clinical evidence



## 1. Introduction

Partial knee arthroplasty (PKA), also termed unicompartmental knee arthroplasty (UKA) when associated with a single compartment, has been performed for isolated single compartment knee osteoarthritis (OA) since the 1970s.<sup>1</sup> PKA can be carried out in the medial, lateral and/or patellofemoral (PF) compartments.

When compared to total knee arthroplasty (TKA), studies have shown that medial PKA patients experience greater retention of normal knee kinematics and accelerated recovery, while suffering less blood loss and reduced postoperative morbidity.<sup>2-5</sup> Lateral PKA is less common, comprising around one-eighth of all PKA cases.<sup>6</sup> However, lateral PKA has also been shown to be an effective treatment in the appropriate patient, with survivorship and outcomes comparable to medial PKA.<sup>6-8</sup> PF arthroplasty has also demonstrated significant benefits to the patient when compared to TKA. A 2018 double-blinded study showed that patients who underwent PF arthroplasty for isolated PF arthritis had a better overall knee-specific quality of life than patients who underwent TKA throughout the first two years after the operation.<sup>9</sup>

Despite the volume of evidence demonstrating the benefits of PKA, the procedure is known to be sensitive to surgical factors such as implant positioning and soft tissue balance.<sup>10</sup> This was recently highlighted in a study by Kazarian et al.,<sup>11</sup> where data from 253 medial PKA patients was retrospectively analyzed to assess the implant survival and radiographic outcomes after PKA, as well as the impact of component alignment and overhang on implant survival. All procedures in the study were performed by two high-volume surgeons. The results showed that the incidence of PKA revisions and alignment outliers were greater than expected, even among high-volume arthroplasty surgeons performing an average of 14.2 PKAs per year. Both alignment and overhang outliers were significant risk factors for implant failure.<sup>11</sup> The researchers emphasize that the ability of low-volume PKA surgeons to consistently attain accuracy in implant position is an important factor to investigate to help enhance PKA survivorship.<sup>11</sup> The Mako System was introduced to provide accurate implant alignment and anatomic restoration and soft-tissue balancing, thereby helping the surgeon restore native knee kinematics and enhance patient outcomes.<sup>12-14</sup> This document summarizes the evidence to date that supports the use of Mako Robotic-Arm Assisted Surgery for PKA (Mako Partial Knee).


## 2. What evidence is available on Mako Partial Knee?

Successful clinical outcomes following joint replacement are dependent on component placement and on restoring the natural kinematics of the knee. Component malalignment in UKA has been associated with stress concentrations, bone fracture and poor clinical outcomes.<sup>15,16</sup> The Mako System is designed to minimize the margin of error associated with component placement and to enhance the accuracy and reproducibility of PKA. Additionally, the Mako System helps enable the surgeon to dynamically balance soft tissue tensioning intraoperatively, with the goal of recreating natural knee kinematics. Clinical studies have shown that Mako Partial Knee has the potential to produce accurate and reproducible component placement in accordance with preoperative plans<sup>17</sup> and to reestablish soft tissue balance.<sup>18</sup>

### 2.1 Component placement accuracy

A key clinical paper on Mako accuracy, published by Bell et al., reports on a randomized controlled trial (RCT) involving 120 patients. The study compared patients who received robotic-arm assisted PKA (Restoris MCK n=62) with those who underwent manually implanted PKA (Oxford n=58).<sup>17</sup> Comparisons were made between groups in terms of the preoperative plan of femoral and tibial component positioning against the actual alignment achieved in three different planes (axial, coronal and sagittal). Results showed more accurate component positioning in the robotic-arm assisted group, with lower root mean square (RMS) errors and significantly lower median errors in all six component parameters ( $p < 0.01$ ).<sup>17</sup> The proportion of patients with tibial slope within 2° of the target position was significantly greater using the robotic-

**Percentage of knees with components positioned within 2° of the target value**



**Figure 1. Bell et al. (2016) showed that use of robotic-arm assisted PKA enabled surgeons to place the tibial and femoral components more accurately and consistently to plan. FS= Femoral Sagittal, FC= Femoral Coronal, FA= Femoral Axial, TS= Tibial Sagittal, TC\*= Tibial Coronal, TA= Tibial Axial. \* = non-significant parameter.<sup>17</sup>**


arm assisted technique than the manual technique (80% compared with 22%,  $p=0.0001$ ). It was concluded that the Mako System more consistently placed the PKA implant in accordance with the preoperative plan (Figure 1).<sup>17</sup>

These results were corroborated by a 2018 study performed at University College Hospital in London, England, by Kayani et al.<sup>19</sup> A single surgeon compared implant placement accuracy using radiographs from 60 consecutive conventional PKAs (Oxford) compared to the surgeon's first 60 consecutive Mako Partial Knees (Restoris MCK). The Mako group had significantly ( $p<0.001$ ) more accurate placement to plan for the femoral and tibial implants, as well as more accurate recreation of the knee's mechanical alignment, posterior tibial slope and joint line height.<sup>19</sup>

## 2.2 Surgical team learning curve

During this initial set of 60 Mako Partial Knee cases within the Kayani et al. study, the surgeon also noted a learning curve of six cases for operating time and surgical team confidence levels to become consistent with conventional PKA statistics.<sup>19</sup> The learning curve did not influence any of the associated accuracy variables, and accuracy to plan achieved with the Mako System was consistent between the surgeon's first Mako case and last 10 Mako cases. This indicated that Mako Partial Knee surgery did not have a learning curve for accuracy in achieving the planned femoral and tibial implant position. Further, no additional risk was observed for postoperative complications during the surgical team learning curve.<sup>19</sup>

### Learning curve



**Figure 2. Typical Mako Partial Knee learning curve graph showing one surgeon's first 50 cases from a multicenter study by Jinnah et al. (2010). After approximately 13 cases, surgical time reached a steady state.<sup>20</sup>**

Jinnah et al. have previously performed an extensive multicenter study to understand how learning curve may influence surgical time for Mako Partial Knee.<sup>20</sup> Eight hundred and ninety-two patients had a Mako Partial Knee performed by 13 different surgeons. Surgical time was measured from insertion of the first bone pin to the acceptance of the final trial components. The average surgical time for all surgeons was  $56 \pm 20$  minutes. The shortest average surgical time for an individual surgeon was  $38 \pm 9$  minutes and the longest was  $70 \pm 29$  minutes. An average learning curve of 13 cases was proposed for the surgical time to reach a steady state (Figure 2).<sup>20</sup>

## 2.3 Soft tissue balance and bone preservation

From a soft tissue perspective, Plate et al. considered that the ability to effectively restore a patient's ligament length and tension may help with restoration of normal knee kinematics and muscle lever arms of the knee joint.<sup>18</sup> Their study examined the accuracy of dynamic, real-time ligament balancing for 52 Mako Partial Knees. Gap distances at  $0^\circ$ ,  $30^\circ$ ,  $60^\circ$ ,  $90^\circ$  and  $110^\circ$  flexion were assessed preoperatively and after final component implantation to establish whether ligament balancing was restored. Ligament balancing was accurate up to 0.53 mm compared to the preoperative plan.<sup>18</sup> These results indicated the Mako System was capable of accurately and precisely reproducing the desired soft tissue balance.

In addition to this, a cadaveric investigation was carried out with the aim of quantifying the amount of bone preserved in robotic medial PKA compared to robotic TKA.<sup>21</sup> Eleven knees were selected and analyzed from seven cadavers. Results showed that robotic PKA procedures resected an average of  $11.6 \pm 1.33$  cm<sup>3</sup> (range: 9.85-13.7 cm<sup>3</sup>) whereas total knee procedures resected an average of  $59.7 \pm 9.65$  cm<sup>3</sup> (range: 47.4-78.3 cm<sup>3</sup>), demonstrating that for this study population, only 17% to 19% of the bone volume was resected in robotic PKA compared to robotic TKA. The study highlighted that in robotic PKA, the femur preparation is contoured to match the implant, which may in turn contribute to enhanced bone preservation and retention of bone stock.<sup>21</sup>

## 2.4 Summary of evidence

These studies demonstrated that robotic-arm assisted technology equipped the surgeon to accurately and consistently place the femoral and tibial PKA components<sup>17</sup> in accordance with preoperative plans and to effectively restore soft tissue balancing.<sup>18</sup> This technology is associated with a short learning curve to achieve time neutrality compared to manual surgery, without influencing the ability to achieve high accuracy.<sup>19</sup>

### 3. What are the potential clinical benefits of Mako Partial Knee?

Mako Partial Knee has been shown to deliver demonstrable clinical benefits.<sup>12-14,24-32</sup> Studies have investigated implant survivorship, patient satisfaction, clinical outcomes and functional outcomes in medial Mako Partial Knee, with favorable results in comparison to other surgical methods.<sup>12-14,24-32,55</sup> In lateral and PF Mako Partial Knee, promising clinical and functional outcomes have also been observed.<sup>35-37,42</sup> Furthermore, in both medial and lateral PKA, congruence of the nonsurgical and surgical compartments has been found to be restored, supporting the hypothesis that the resultant redistribution of contact forces across the patellofemoral joint could help address PF symptoms.<sup>35-37,42</sup>

#### 3.1 Survivorship

Favorable survivorship data was published in the U.S. by Burger et al.<sup>30</sup> in a study that evaluated midterm implant survivorship for robotic-arm assisted PKA patients.<sup>30</sup> The research involved a retrospective review of patients who underwent robotic-arm assisted PKA between 2007 and 2016. Study participants received a fixed-bearing medial or lateral PKA, patellofemoral arthroplasty (PFA), or bicompartamental knee arthroplasty (involving a PFA plus medial PKA), and the mean follow-up was 4.7 years (2.0 to 10.8).<sup>30</sup> The five-year survivorship rate of medial PKA (n=802), lateral PKA (n=171) and PFA/ bicompartamental knee arthroplasty (n=35/10) was 97.8%, 97.7% and 93.3%, respectively.<sup>30</sup>

Comparable data was previously confirmed in a multicenter longitudinal study evaluating short- and midterm survivorship of robotic-arm assisted medial PKA, which demonstrated 98.8% survivorship (in 909 knees) at 2.5-year follow-up and 97% survivorship (in 432 knees) at 5.5-year follow-up.<sup>12,22</sup> This survivorship rate was greater than rates derived from high-volume surgeon data and registry data for conventional PKA (Figure 3).<sup>12,22</sup> The study concluded that the favorable survivorship observed resulted from Mako's ability to help enable surgeons to achieve more accurate component positioning when compared to implant placement using manual techniques.<sup>12,22</sup>

An RCT by Gilmour et al., comparing patients who underwent medial Mako Partial Knee (Restoris MCK) with those who underwent manual medial PKA (Oxford) demonstrated encouraging results.




Figure 3. Survivorship data from Pearle et al. (2017)<sup>22</sup> and Kleebblad et al. (2018)<sup>12</sup> on robotic-arm assisted PKA compared to studies in literature and annual registries reporting 2 to 3 years and 5 to 6 years conventional PKA survivorship data.




Figure 4. Graph indicating Mako Partial Knee revision rates with data taken from cohort studies, economic analyses, level I clinical trials (RCTs) and international registries.<sup>12, 22, 24, 30, 46, 48-51, 54</sup>

Specifically, Mako Partial Knee patients had 100% survivorship compared to 96.3% in the manual group at two years postoperation.<sup>23</sup> The 100% survivorship rate was maintained in the robotic group at five years postoperation.<sup>49</sup>

Similar promising data was published in the 2019 Australian Joint Registry,<sup>24</sup> which reported the cumulative percent revision for Restoris MCK medial PKA as 1.5% at one year and 2.5% at three years. These rates were much lower than all non-robotic PKAs, which were reported to be 2.1% and 4.6% at one and three years, respectively. The cumulative revision rate of the Restoris MCK medial PKA also compared favorably to the revision rate for all Oxford medial PKA replacements, which were 2.2% at one year and 5.8% at three years.<sup>24</sup> These findings were reflected in a study conducted by St Mart et al.,<sup>55</sup> who examined the cumulative revision rate of PKA procedures implanted with the Mako System using data from the Australian Joint Registry between 2015 and 2018. The researchers found that the Mako-assisted Restoris had significantly lower overall revision rate compared to other types of non-robotically assisted PKA procedures. However, the higher rate of early revision for infection for robotically assisted PKA requires further investigation.<sup>55</sup>

The revision rates for Mako Partial Knee have been published in cohort studies, economic analyses, level I clinical trials (RCTs) and international registries (Figure 4). The evidence supports excellent survivorship of the Restoris MCK implant.

### 3.2 Patient satisfaction

In a multicenter longitudinal clinical trial, the vast majority of Mako Partial Knee patients were “very satisfied” or “satisfied” with their joint replacement.<sup>12,22</sup> This study performed follow-up at 2.5 years (909 knees) and 5.5 years (432 knees) with patients who underwent medial Mako Partial Knee procedures.<sup>12,22</sup> 92% of patients reported




Figure 5. Midterm patient satisfaction with medial Mako Partial Knee procedures (Kleebblad et al., 2018<sup>12</sup> and Pearle et al., 2017<sup>22</sup>).

satisfaction with their knee 2.5 years postoperatively and 91% of patients reported satisfaction at 5.5 years (Figure 5).<sup>12,22</sup> In a similar study based on the Swedish Knee Arthroplasty Registry, 83% of 7,860 patients who underwent manual medial PKA were satisfied with their knee at an average six-year follow-up.<sup>25</sup>

Using the Mako System, Coon et al. performed 152 (71.3%) medial PKAs, 33 (15.5%) lateral PKAs, 20 (9.4%) medial bicompartamental PKAs and 8 (3.8%) patellofemoral PKAs. All surgical procedures had high patient satisfaction with an average of 82.5% of patients reporting being very satisfied or satisfied at six months, which increased to 89.5% at two years.<sup>28</sup> The lateral PKA group reported 100% satisfaction two years postoperation.<sup>29</sup>

Comparable midterm patient satisfaction data was recently published in a large single-surgeon study of 1018 knees, where a large proportion of patients who underwent robotic-assisted PKA reported high satisfaction levels.<sup>30</sup> The mean follow-up was 4.7 years (2.0 to 8.0), and the results showed that 90.7% of medial PKA patients, 92.6% of lateral PKA patients and 78.9% of PFA or bicompartamental knee arthroplasty patients were either very satisfied or satisfied with their knee function.<sup>30</sup>

### 3.3 Clinical outcomes

An RCT performed by Blyth et al. found that patients who underwent medial Mako Partial Knee experienced less pain than those who underwent manual surgery during the 90-day postoperative period.<sup>13</sup> Median pain scores were 55.4% lower in robotic-arm assisted patients compared to manual patients from day one to day 56 (Figure 6).<sup>13</sup> Furthermore, the robotic-arm assisted patients had a better American Knee Society Score (AKSS) at three months postoperatively and at one year postoperatively, and a greater proportion of robotic-arm assisted patients




Figure 6. Visual analog pain score collected up to 90 days postoperatively in a RCT of manual vs. robotic arm-assisted medial PKA procedures.<sup>13</sup>

showed improvements in their UCLA Activity Score.<sup>13</sup> Through binary logistic regression, the study was also able to predict the key factors associated with achieving excellent outcomes on the AKSS. These factors were a preoperative UCLA Activity Score level >5 and the use of robotic-arm assisted surgery, although these do not withstand adjustment for multiple comparisons.<sup>13</sup>

In two separate studies, evidence showed that medial Mako Partial Knee patients were more likely to “forget” their artificial joint during daily life compared to those who underwent manual TKA.<sup>26,27</sup> Zuiderbaan et al. administered The Forgotten Joint Score (FJS) questionnaire one and two years postoperatively.<sup>26</sup> Scores were compared between 65 patients who underwent medial Mako Partial Knee and 65 patients who underwent manually instrumented TKA.<sup>26</sup> Results demonstrated patients who underwent medial robotic-arm assisted PKA were more likely to forget their artificial joint in daily life (Figure 7).<sup>26</sup> Similarly, in a separate powered cohort study from the U.K., conducted by Clement et al.<sup>27</sup>, 30 patients who underwent Mako PKA were propensity score matched to 90 patients who underwent manual TKA for isolated medial compartment arthritis. The findings from this study showed that the six-month FJS was significantly greater for the robotic PKA group compared to the manual TKA group (difference 24.2,  $p < 0.001$ ) (Figure 7).<sup>27</sup>

Furthermore, the same powered (1:3 ratio) cohort study by Clement et al. published encouraging early postoperative outcomes data, where statistically and clinically significant greater knee-specific functional outcomes were observed in robotic PKA patients compared to those who underwent manual TKA.<sup>27</sup> Findings showed that the robotic PKA group had a significantly greater six-month Oxford Knee Score by nearly eight points, and there was a five-point (95% confidence interval 1.9 to 8.1;  $p < 0.001$ ) greater improvement in the robotic PKA group compared to the manual TKA group, which was greater than the




Figure 7. FJS at six months, one and two years post-operation showing significantly higher scores in the medial Mako Partial Knee group ( $p < 0.001$ ,  $p = 0.002$  and  $p = 0.004$ , respectively)<sup>26,27</sup>

minimal clinically important difference. This positive early outcome data was further fortified within the study as the researchers also found that the robotic PKA group had significantly better postoperative pain visual analog scale (VAS) scores compared with the manual TKA group (Table 1).<sup>27</sup>

Table 1. Six-month postoperative outcome measures and differences between robotic PKA vs. manual TKA.<sup>27</sup>

| Mean PROM (SD)         | rUKA          | mTKA          | Difference (95% CI)    | p-value* |
|------------------------|---------------|---------------|------------------------|----------|
| Postoperative OKS      | 44.2 (4.4)    | 36.5 (9.4)    | 7.7 (4.2 to 11.3)      | <0.001   |
| Postoperative FJS      | 77.1 (25.9)   | 52.9 (32.6)   | 24.2 (11.2 to 37.2)    | <0.001   |
| Postoperative EQ-5D    | 0.913 (0.126) | 0.764 (0.248) | 0.148 (0.054 to 0.241) | 0.002    |
| Postoperative pain VAS | 93.6 (12.3)   | 76.4 (24.8)   | 20.5 (9.9 to 31.0)     | <0.001   |

\*Unpaired t-test. CI, confidence interval; EQ-5D, EuroQol five-dimension questionnaire; FJS, Forgotten Joint Score; mTKA, manual total knee arthroplasty; OKS, Oxford Knee Score; PROM, patient-reported outcome measure; rUKA, robotic unicompartmental knee arthroplasty; VAS, visual analog scale.

Overall, results of these studies suggested positive clinical and patient-reported outcomes of robotic-arm assisted medial, lateral, PF and bicompartmental PKA.<sup>10,12-14,20,23,27-29,30,55</sup>

### 3.4 Functional outcomes

Gait analysis has been used to compare outcomes of robotic-arm assisted PKA patients to those of manual Oxford PKA patients. In an RCT, Motesharei et al. compared the gait of 31 robotic PKA patients to 39 Oxford PKA patients one year postoperatively.<sup>31</sup> Both groups were compared to a control group of 50 healthy subjects obtained from the University of Strathclyde’s archive.<sup>31</sup> Results from this study showed statistically significant differences in knee joint kinematics during level walking between the robotic-arm assisted and manual PKA groups.




Figure 8. Mean knee excursion angles of the control group, the robotic-arm assisted and manual PKA groups during the stance phase of gait at one year post-operation.<sup>31</sup>

The robotic-arm assisted group achieved a higher knee excursion (18.0°, SD 4.9°) compared to the manual group (15.7°, SD 4.1°) (Figure 8 and Table 2).<sup>31</sup> There was no significant difference between the healthy group and the robotic-arm assisted group, but there was a significant difference between the healthy group and the manual group (p<0.001).<sup>31</sup>

This study was repeated at five-years postoperatively by Millar et al., though on fewer patients (25 Mako vs. 21 Oxford), and the differences seen at one year were maintained.<sup>14</sup> Results showed that the Mako group achieved significantly greater knee flexion in weight acceptance than the conventional group (Table 3).<sup>14</sup> These findings suggested that the improved alignment offered by the Mako System may result in enhanced function of the knee during gait, and that the use of the Mako System resulted in a gait pattern that facilitated the normal function of the knee more closely than the conventional technique.<sup>14,31</sup>

A clinical study by Borus et al. assessed functional performance in patients who received robotic-arm assisted PKA compared to those who received manual TKA.<sup>32</sup> Tests included a six-minute walk, timed up and go, and stair ascend/descend, which were measured preoperatively and at six weeks and at three months postoperatively. Although a statistically significant difference in functional performance change between groups was not reached, the authors highlighted that at six weeks, the robotic PKA group was able to walk an additional 21.00 meters (68.90 feet) compared to just 5.95 meters (19.52 feet) for the manual TKA group.<sup>32</sup> Very similar functional differences were observed with the timed up and go and stair ascend/descend tests, suggesting that robotic PKA provided functional benefits that were at least equivalent to manual TKA.<sup>32</sup>

Research by Coon et al. on medial Mako Partial Knees, lateral PKAs, medial bicompartamental PKAs and patellofemoral PKAs showed that at two years postoperatively, 87.9% of patients were as active or the

**Table 2. Comparison of knee excursion values during loading phase of gait at one year post-operation. Standard deviation in brackets.**<sup>31</sup>

|  | Control  | Mako       | Oxford      |
|--|--|------------|-------------|
| <b>Knee excursion from foot-strike to mid-stance (degrees)</b> | 19.5 (4.0)                                       | 18.0 (4.9) | 15.7 (4.1)* |
|  | * Significantly different than the control group |            |             |

**Table 3. Mean (SD) excursion during weight acceptance for each patient group at five years post-operation.**<sup>14</sup>

| Patient group | Mean (SD) excursion during WA (°) |
|---------------|-----------------------------------|
| <b>Mako</b>   | 14.3 (6.4)                        |
| <b>Oxford</b> | 9.9 (4.2)                         |
| <b>P</b>      | 0.008                             |


same as they expected they would be before surgery.<sup>29</sup> In addition, the average distance walked at discharge was 79.8 meters, and 90.9% of patients were walking without support three weeks postoperatively.<sup>29</sup> Lastly, 65 patients were employed at time of surgery, and 86% of those patients returned to work six weeks after their operation.<sup>29</sup>

### 3.5 Clinical outcomes of lateral PKA

Lateral PKA is less frequently performed within the general population, accounting for just one-eighth of PKA cases.<sup>6</sup> However, this procedure has been shown to be effective for the appropriate patient, achieving reliable improvements in pain, function and implant survivorship.<sup>6-8</sup> The Mako robotic platform offers potential benefits through its demonstrated accuracy and reproducible implant positioning, helping to minimize the margin of error associated with component placement.<sup>12</sup> In addition, the platform enables intraoperative dynamic soft tissue balancing to help the surgeon recreate the patient's natural knee kinematics.

Promising results have been reported by several studies examining lateral Mako Partial Knee.<sup>33,34</sup> For example, a retrospective study conducted by van der List et al. compared two-year postoperative functional outcomes using the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC score) and FJS, between 143 medial and 36 lateral Mako Partial Knee procedures (Figure 9, Table 4).<sup>33</sup> Equivalent functional outcomes were noted for both medial and lateral PKA procedures.<sup>33</sup>

Similar promising survivorship data was published by Augart et al.<sup>34</sup> The authors performed a search of their institution's joint registry and found 88 lateral robotic-arm assisted PKA patients, with a mean follow-up of 24.4 months ± 10.7 months, who had 100% survivorship at final follow-up without revision to TKA.<sup>34</sup> The promising data observed thus far from medial and lateral Mako Partial Knees suggests that the potential benefits offered by the Mako robotic platform, with regards to surgical planning, precision, reproducibility and intraoperative soft tissue adjustments, have the potential to help enhance surgical accuracy during these technically demanding procedures.



**Figure 9. Preoperative and postoperative radiographs of: medial Mako Partial Knee (left), and lateral Mako Partial Knee (right).**<sup>33</sup>

**Table 4. Mean ( $\pm$ SD) scores of WOMAC and FJS of all patients undergoing medial and lateral UKA and stratified by postoperative alignment as neutral or undercorrected.<sup>33</sup>**


| Postoperative alignment                              | Score | N   | Medial UKA      | N  | Lateral UKA     | Medial vs. lateral |
|--|-------|-----|-----------------|----|-----------------|--------------------|
| All patients <sup>a</sup>                            | WOMAC | 143 | 89.8 $\pm$ 11.7 | 36 | 90.2 $\pm$ 12.4 | 0.855              |
|  | FJS   | 95  | 71.2 $\pm$ 24.5 | 25 | 70.9 $\pm$ 28.2 | 0.956              |
| Neutral aligned patients ( $-1^\circ$ to $3^\circ$ ) | WOMAC | 85  | 90.9 $\pm$ 11.4 | 19 | 87.2 $\pm$ 12.5 | 0.200              |
|  | FJS   | 57  | 72.6 $\pm$ 22.6 | 12 | 55.3 $\pm$ 28.5 | 0.024*             |
| Undercorrected patients ( $3^\circ$ to $7^\circ$ )   | WOMAC | 51  | 88.5 $\pm$ 11.6 | 15 | 96.0 $\pm$ 5.4  | 0.001*             |
|  | FJS   | 38  | 68.2 $\pm$ 26.8 | 13 | 85.3 $\pm$ 19.5 | 0.020*             |
| Neutral vs. undercorrected                           | WOMAC | 143 | 0.214           |    | 0.005*          |                    |
|  | FJS   | 143 | 0.199           |    | 0.010*          |                    |

UKA indicates unicompartmental knee arthroplasty; FU, follow-up; WOMAC, Western Ontario and McMaster Universities Arthritis Index; FJS, Forgotten Joint Score. Neutral alignment for medial UKA indicates one degree of valgus to three degrees of varus and for lateral UKA indicates one degree of varus to three degrees of valgus.


Undercorrected alignment for medial UKA indicates three degrees to seven degrees of varus and for lateral UKA indicates three degrees to seven degrees of valgus.

\* Indicates a significant difference with  $p < 0.05$ .

<sup>a</sup> 12 patients with medial UKA and 2 patients with lateral UKA had no postoperative hip-knee-ankle radiograph and could not be included for subgroup analysis.




**Figure 10. Khamaisy et al. (2016). Iterative closest point algorithm was performed to calculate the congruence index (noted as INDX in the figure) of the lateral compartment pre- and postoperatively following manual digitization of the femoral and tibial surfaces in patients who received a medial Mako Partial Knee.<sup>37</sup>**



**Figure 11. Preoperative Merchant view of a left knee. The trochlear angle (red angle) is  $140^\circ$ . The congruence angle (yellow angle) is  $14^\circ$ . The medial patellofemoral joint space is represented by the purple line.<sup>36</sup>**

### 3.6 Continuum of care

As mean patient age decreases, partial knee arthroplasty is often indicated as a conservative treatment to delay need for a total knee replacement. Studies of joint line restoration, patella tracking, and medial and lateral compartment congruency have been conducted at Hospital for Special Surgery in New York.<sup>35-37</sup> In all three studies, congruence of the surgical compartment was restored through the Mako procedure and implant.<sup>35-37</sup> Congruence and joint line of the nonoperative compartment were also restored ( $p=0.001$ ).<sup>35</sup> The authors hypothesized that the improved patellofemoral congruence after Mako Partial Knee may lead to redistribution of contact forces across the patellofemoral joint and secondarily treat PF symptoms (Figure 10, Figure 11, Figure 12).<sup>34,35</sup>



**Figure 12. Postoperative Merchant view of a left knee. The trochlear angle (red angle) is  $140^\circ$ . The postoperative congruence angle (yellow angle:  $6^\circ$ ) was decreased compared to the preoperative one (Figure 11). Moreover, the medial patellofemoral joint space (purple line) was increased by 1.5 mm following PKA.<sup>36</sup>**

Similarly, in a 2020 study by Burger et al.,<sup>38</sup> researchers aimed to explore the effect of patellofemoral joint pathology on lateral PKA. In particular, the effect of preoperative radiological degenerative changes and alignment on patient-reported outcome measures (PROMs) after lateral PKA was evaluated, as well as the influence of lateral PKA on the alignment of the patellofemoral joint.<sup>38</sup> A consecutive series of 140 knees in 130 patients who underwent Mako robotic arm-assisted fixed-bearing lateral PKA were retrospectively reviewed. Radiological evaluation was conducted to obtain a Kellgren Lawrence (KL) grade, an Altman score and alignment measurements for each knee. Postoperative PROMs were assessed using the Kujala (Anterior Knee Pain Scale) score, Knee Injury and Osteoarthritis Outcome Score Joint Replacement (KOOS JR) and satisfaction levels. The results showed that at mean 4.1 years (2.0 to 8.5) follow-up, good to excellent Kujala scores were reported, and the presence of mild to moderate preoperative patellofemoral joint osteoarthritis had no impact on these scores (KL grade 0 vs. 1 to 3,  $p = 0.203$ ; grade 0 to 1 vs. 2 to 3,  $p = 0.674$ ). Comparable scores were reported by patients with osteoarthritis evident on either the medial or lateral patellofemoral joint facet, and patients with abnormal patellar congruence and tilt angles ( $\geq 17^\circ$  and  $\geq 14^\circ$ , respectively) reported good to excellent Kujala scores. Furthermore, it was evident that lateral PKA resulted in improvements to patellofemoral alignment.<sup>38</sup> The findings from this study are encouraging as this is the first study demonstrating that mild to moderate preoperative radiological degenerative changes and malalignment of the patellofemoral joint are not associated with poor patient-reported outcomes at midterm follow-up after lateral fixed-bearing PKA. The researchers went on to suggest that this may be explained by realignment of the patella and the resulting redistribution of loads across the patellofemoral joint.<sup>38</sup>

### 3.7 Outcomes of patellofemoral arthroplasty

The purpose of patellofemoral arthroplasty is to address the pain caused at the patellofemoral joint without performing a more substantial total knee surgery that would sacrifice additional bone. However, past literature has reported conflicting success rates of PFA as a surgical treatment for patellofemoral OA.<sup>39-40</sup> Odgaard et al. used a multicenter, double-blinded RCT to compare clinical outcomes associated with PFA and TKA to establish whether there was an advantage to either option.<sup>41</sup> They found that PFA patients recovered quicker than TKA patients, and the functional outcomes were also better for PFA patients.<sup>41</sup> The average TKA patient lost almost three months of knee function postoperatively during the first two years, relative to the PFA patient.<sup>41</sup> It was concluded that PFA was a superior option to TKA in patients with patellofemoral OA.<sup>41</sup>

Encouraging functional data was observed in another study by Noyes et al., which examined the early results of 33 prospective, consecutive third-generation Mako PFA procedures.<sup>42</sup> The authors analyzed both sports and work activity levels in younger active patients. The study included 33 consecutive PFAs in 29 patients (four bilateral), with a mean age 40 (range, 22-68).<sup>42</sup> All patients received a comprehensive clinical evaluation, Cincinnati Knee Rating and International Knee Documentation Committee (IKDC) objective rating. They also received radiographic evaluation. Results showed high levels of participation in light sports: 22% preoperatively, rising to 87% postoperatively. A total of 85% of patients in the employed subgroup returned to work postoperatively, and in six out of seven patients who received surgery due to articular cartilage restoration failure, improvement was seen postoperatively and they returned to light sports/work.<sup>42</sup> This research demonstrated that robotic-arm assisted PFA was a successful treatment option in younger active patients with isolated PF arthritis, enabling the majority of those patients to return to low-impact recreational activities and occupations.<sup>42</sup>

#### 4. Is Mako cost-effective?

Compared to TKA, studies have shown that UKA patients have fewer postoperative complications,<sup>43</sup> improved FJS,<sup>26,27</sup> and higher quality-adjusted life-years (QALYs) in older patients.<sup>52-54</sup>

With rising demand for PKA in patients who seek restored function and a quicker recovery time, a U.S. study performed by Kazarian et al. evaluated the cost-effectiveness of PKA compared to TKA as well as nonsurgical treatment (NST).<sup>52</sup> Using a Markov decision analytic model, the authors assessed lifetime costs and QALYs as function of age at time of initial treatment (ATIT) of patients with end-stage unicompartmental knee osteoarthritis. The analysis included direct medical and indirect costs. Models were run for ATITs at five-year intervals from 40 through 90 years of age. Results indicated PKA had the greatest QALY accumulation followed by TKA and NST, and that PKA was more cost-effective compared to NST for patients aged 40 to 86. Furthermore, when surgical treatments were compared, PKA dominated TKA by generating more QALYs than TKA for all ATITs. The authors further concluded that if PKAs were performed as 12% to 20% of the total volume of knee arthroplasties versus the less than 8% observed, it would lead to a lifetime cost-savings of 987 million to 1.5 billion U.S. dollars and increased lifetime QALY accumulation of 124,403 to 217,705 across the U.S. population.<sup>52</sup>




Figure 13. ICER values comparing PKA with NST, TKA with NST, and PKA with TKA by age.<sup>52</sup>

In a separate U.K.-based study, a Markov decision analysis was performed to assess the cost-effectiveness of robotic PKA (rPKA) relative to manual TKA (mTKA) and manual PKA (mPKA) for patients with isolated medial compartment OA of the knee with a mean age of 65 years.<sup>53</sup> The study objective was to identify the cost per quality adjusted life-year of rPKA relative to mTKA and mPKA. Model inputs included hospital costs, implant survival and mortality rate. Using a model with an annual case volume of 100 patients, the cost per QALY of rPKA was £1,395 and £1,170 relative to mTKA and mPKA, respectively. The cost per QALY was influenced by case volume: a low-volume center performing 10 cases per year would achieve a cost per QALY of £7,170 and £8,604 relative to TKA and PKA, respectively. For a high-volume center performing 200 rPKAs per year with a mean two-day length of stay, the cost per QALY would be £648; if performed as day cases, the cost would be reduced to £364 relative to TKA. For a high-volume center performing 200 rPKAs per year with a shorter length of stay of one day relative to PKA, the cost per QALY would be £574 (Figure 14<sup>53</sup>). Furthermore, the cost per QALY of rPKA decreased with reducing length of hospital stay and with increasing case volume, compared with mTKA and mPKA.<sup>53</sup> The model showed that rPKA was a cost-effective alternative to mTKA and mPKA for patients with isolated medial compartment OA of the knee.




Figure 14. Cost per quality-adjusted life-year (QALY) or robotic-assisted PKA according to case volume and length of hospital stay relative to TKA.<sup>53</sup>

In summary, these models demonstrated that in patients with isolated medial compartment arthritis, PKA was observed to be a more cost-effective procedure compared to nonsurgical treatment and TKA for the specified age groups modeled, thus concluding rPKA was cost-effective compared to TKA.

A hospital in Brisbane, Australia examined the potential cost-savings for the health system and the community in a broadly accessible model through the increased utilization of PKA using robotic-arm assisted PKA vs. conventional TKA.<sup>42</sup> They retrospectively reviewed 240 patients where the first 120 consecutive Mako Partial Knees performed during this period were matched to 120 conventional TKAs. Clinical data from the medical records and costs for procedure for each component were collected. Bivariate analyses were performed on the data to determine if there were statistically significant differences by surgery type in clinical outcomes and financial costs. The study found a significantly lower cost incurred for robotic-arm assisted PKA vs. TKA with an average savings of AU\$7,179 per case. The operating time (86.0 min vs. 75.9 min;  $p=0.004$ ) was significantly higher for PKA but the length of stay was significantly lower (1.8 vs. 4.8 days;  $p<0.001$ ). This study also found a significant difference in the use of opioids in PKA compared to TKA (125.0 morphine equivalent (ME) vs. 522.1 ME,  $p<0.001$ ).<sup>42</sup>

In the U.S., in a study by Cool et al., reasons for revisions and associated costs were analyzed for unicompartmental arthroplasty cases.<sup>54</sup> UKA procedures were identified using a commercial administrative claims database to evaluate hospital admissions for revision surgeries. Robotic UKA (rUKA, Mako Partial Knee) and manual UKA (mUKA, manual partial knee) procedures performed between March 1, 2013 and July 31, 2015 were used to calculate the rate of revisions within 24 months of the index procedure. Cases were propensity matched 2:1 based on age, sex, race, geographic division, high-cost comorbidities and concentration of healthcare specialists per 100,000 population to control for outside confounding factors at case index. A total of 738 commercial health plan patients (246 rUKA, 492 mUKA) were selected for inclusion in the analysis. Results indicated fewer revision procedures in rUKA (0.81% (2/246) vs. 5.28% (26/492);  $p=0.0017$ ) and rUKA patients incurred lower mean costs for the index stay plus revision(s) (\$26,001 vs. \$27,977;  $p>0.05$ ). Lower length of stay at index was also noted in the rUKA group (1.77 vs. 2.02 days;  $p=0.0047$ ). The study concluded that patients who underwent rUKA had fewer revision procedures, shorter LOS and incurred lower mean costs at 24 months.<sup>54</sup>

Findings from a 2020 U.K. cohort study involving 30 Mako Partial Knees compared to 90 propensity-matched manual TKAs showed that the length of stay was significantly ( $p < 0.001$ ) shorter in the robotic-arm assisted PKA group (median two days, interquartile range (IQR) one to three) compared to the manual TKA group (median four days, IQR three to five). The shorter length of stay observed in this study was considered a cost saving for the center relative to mTKA.<sup>27</sup>

The cost-effectiveness studies described above all differed in inputs specific to their country, local region, hospital system or payer. These studies demonstrated that robotic-arm assisted partial knee arthroplasty, in comparison to manual TKA or manual partial knees, was associated with lower costs and/or improvements in QALY.<sup>45,52,53</sup>

## 5. Conclusion

Mako Partial Knee offers the potential for surgeons to achieve component placement accuracy<sup>17</sup> and soft tissue balancing,<sup>18</sup> as well as to enhance clinical outcomes.<sup>12-14,22-32</sup> Patients have reported tangible benefits of robotic-arm assisted procedures, including treatment satisfaction,<sup>12,22,28,26,30</sup> return to activities of daily living<sup>28</sup> and a “forgotten” joint.<sup>13,26,27</sup> Surgeons are empowered to achieve their target preoperative plans with precision,<sup>17</sup> helping distinguish them within their medical communities. The cost-effectiveness studies described here demonstrated favorable economic returns, lower costs and better improvements in QALY for patients who received robotic-arm assisted partial knees in contrast to those received TKA or manual partial knees.<sup>45,52,53</sup> Ultimately, the benefits of Mako Partial Knee surgery are reported to be experienced by all key players – patients, surgeons and health systems.

**Notes**

**Notes**

## References

- Ollivier M, Abdel MP, Parratte S, Argenson JN. Lateral unicompartmental knee arthroplasty (UKA): contemporary indications, surgical technique, and results. *Int Orthop*. 2014;38(2):449-455. doi:10.1007/s00264-013-2222-9
- Price AJ, Rees JL, Beard DJ, Gill RH, Dodd CA, Murray DM. Sagittal plane kinematics of a mobile-bearing unicompartmental knee arthroplasty at 10 years: a comparative in vivo fluoroscopic analysis. *J Arthroplasty*. 2004;19(5):590-597. doi:10.1016/j.arth.2003.12.082
- Schwab PE, Lavand'homme P, Yombi JC, Thienpont E. Lower blood loss after unicompartmental than total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc*. 2015;23(12):3494-3500. doi:10.1007/s00167-014-3188-x
- Brown NM, Sheth NP, Davis K, et al. Total knee arthroplasty has higher postoperative morbidity than unicompartmental knee arthroplasty: a multicenter analysis. *J Arthroplasty*. 2012;27(8 Suppl):86-90. doi:10.1016/j.arth.2012.03.022
- Larsen K, Sorensen OG, Hansen TB, Thomsen PB, Soballe K. Accelerated perioperative care and rehabilitation intervention for hip and knee replacement is effective: a randomized clinical trial involving 87 patients with 3 months of follow-up. *Acta Orthop*. 2008;79(2):149-159. doi:10.1080/17453670710014923
- Jamali AA, Scott RD. Lateral unicompartmental knee arthroplasty. *Tech Knee Surg*. 2005;4(2):79-88.
- Pennington DW, Swienczkowski JJ, Lutes WB, Drake GN. Lateral unicompartmental knee arthroplasty: survivorship and technical considerations at an average follow-up of 12.4 years. *J Arthroplasty*. 2006;21(1):13-17. doi:10.1016/j.arth.2004.11.021
- Volpi P, Marinoni L, Bait C, Galli M, Denti M. Lateral unicompartmental knee arthroplasty: indications, technique and short-medium term results. *Knee Surg Sports Traumatol Arthrosc*. 2007;15(8):1028-1034. doi:10.1007/s00167-007-0342-8
- Odgaard A, Madsen F, Kristensen PW, Kappel A, Fabrin J. The Mark Coventry Award: Patellofemoral Arthroplasty Results in Better Range of Movement and Early Patient-reported Outcomes Than TKA. *Clin Orthop Relat Res*. 2018;476(1):87-100. doi:10.1007/s11999-00000000000000017
- Citak M, Dersch K, Kamath AF, Haasper C, Gehrke T, Kendoff D. Common causes of failed unicompartmental knee arthroplasty: a single-centre analysis of four hundred and seventy one cases. *Int Orthop*. 2014;38(5):961-965. doi:10.1007/s00264-013-2263-0
- Kazarian GS, Barrack TN, Okafor L, Barrack RL, Nunley RM, Lawrie CM. High prevalence of radiographic outliers and revisions with unicompartmental knee arthroplasty. *J Bone Joint Surg Am*. Published online May 8, 2020. doi:10.2106/JBJS.19.01277. doi:10.2106/JBJS.19.01277
- Kleebad LJ, Borus TA, Coon TM, Douchis J, Nguyen JT, Pearle AD. Midterm survivorship and patient satisfaction of robotic-arm-assisted medial unicompartmental knee arthroplasty: a multicenter study. *J Arthroplasty*. 2018;33(6):1719-1726. doi:10.1016/j.arth.2018.01.036
- Blyth MJG, Anthony I, Rowe P, Banger MS, MacLean A, Jones B. Robotic arm-assisted versus conventional unicompartmental knee arthroplasty: Exploratory secondary analysis of a randomised controlled trial. *Bone Joint Res*. 2017;6(11):631-639. doi:10.1302/2046-3758.611.BJR-2017-0060.R1
- Millar LJ, Banger M, Rowe PJ, Blyth M, Jones B, Maclean A. A five-year follow up of gait in robotic assisted vs conventional unicompartmental knee arthroplasty. *Gait Posture*. 2018;65(Supplement 1):31-32. doi:10.1016/j.gaitpost.2018.06.035
- Tsai TY, Dimitriou D, Liow MH, Rubash HE, Li G, Kwon YM. Three-dimensional imaging analysis of unicompartmental knee arthroplasty evaluated in standing position: component alignment and in vivo articular contact. *J Arthroplasty*. 2016;31(5):1096-1101. doi:10.1016/j.arth.2015.11.027
- Aleto TJ, Berend ME, Ritter MA, Faris PM, Meneghini RM. Aleto TJ, Berend ME, Ritter MA, Faris PM, Meneghini RM. Early failure of unicompartmental knee arthroplasty leading to revision. *J Arthroplasty*. 2008;23(2):159-163. doi:10.1016/j.arth.2007.03.020
- Bell SW, Anthony I, Jones B, MacLean A, Rowe P, Blyth M. Improved accuracy of component positioning with robotic-assisted unicompartmental knee arthroplasty: data from a prospective, randomized controlled study. *J Bone Joint Surg Am*. 2016;98(8):627-635. doi:10.2106/JBJS.15.00664
- Plate JF, Mofidi A, Mannava S, et al. Achieving accurate ligament balancing using robotic-assisted unicompartmental knee arthroplasty. *Adv Orthop*. 2013;2013:837167. doi:10.1155/2013/837167
- Kayani B, Konan S, Pietrzak JRT, Huq SS, Tahmassebi J, Haddad FS. The learning curve associated with robotic-arm assisted unicompartmental knee arthroplasty: a prospective cohort study. *Bone Joint J*. 2018;100-B(8):1033-1042. doi:10.1302/0301-620X.100B8.BJJ-2018-0040.R1
- Jinnah R, Lippincott CJ, Horowitz S, Condit MA. The learning curve of robotically-assisted UKA. Presented at: Orthopaedic Research Society (ORS) 56th Annual Meeting; March 6-10, 2010; New Orleans, LA.
- Hampf E, Chang TC, Pearle A. Robotic partial knee arthroplasty demonstrated greater bone preservation compared to robotic total knee arthroplasty. Poster presented at: Orthopaedic Research Society (ORS) Annual Meeting; February 2-5, 2019; Austin, TX.
- Pearle AD, van der List JP, Lee L, Coon TM, Borus TA, Roche MW. Survivorship and patient satisfaction of robotic-assisted medial unicompartmental knee arthroplasty at a minimum two-year follow-up. *Knee*. 2017;24(2):419-428. doi:10.1016/j.knee.2016.12.001
- Gilmour A, MacLean AD, Rowe PJ, et al. Robotic-arm-assisted vs conventional unicompartmental knee arthroplasty: the 2-year clinical outcomes of a randomized controlled trial. *J Arthroplasty*. 2018;33(7S):S109-S115. doi:10.1016/j.arth.2018.02.050
- National Joint Replacement Registry. Hip, Knee & Shoulder Arthroplasty Annual Report 2019. Australian Orthopaedic Association; 2019.
- Robertsson O, Dunbar M, Pehrsson T, Knutson K, Lidgren L. Patient satisfaction after knee arthroplasty: a report on 27,372 knees operated on between 1981 and 1995 in Sweden. *Acta Orthop Scand*. 2000;71(3):262-267. doi:10.1080/000164700317411852
- Zuiderbaan HA, van der List JP, Khamaisy S, et al. Unicompartmental knee arthroplasty versus total knee arthroplasty: Which type of artificial joint do patients forget?. *Knee Surg Sports Traumatol Arthrosc*. 2017;25(3):681-686. doi:10.1007/s00167-015-3868-1
- Clement ND, Bell A, Simpson P, Macpherson G, Patton JT, Hamilton DF. Robotic-assisted unicompartmental knee arthroplasty has a greater early functional outcome when compared to manual total knee arthroplasty for isolated medial compartment arthritis. *Bone Joint Res*. 2020;9(1):15-22. Published 2020 May 16. doi:10.1302/2046-3758.91.BJR-2019-0147.R1
- Coon T, Shi S, DeBattista J. Clinical and functional outcomes of robotic-arm assisted medial unicompartmental knee arthroplasty. Presented at: European Knee Society Arthroplasty Conference; April 20-21, 2017; London, England.
- Coon T, Shi S, DeBattista J, Bhowmik-Stoker M. Clinical and functional outcomes of robotic-arm assisted unicompartmental and bicompartmental knee arthroplasty. Presented at: European Knee Society Arthroplasty Conference; April 20-21, 2017; London, England.
- Burger JA, Kleebad LJ, Laas N, Pearle AD. Mid-term survivorship and patient-reported outcomes of robotic-arm assisted partial knee arthroplasty. *Bone Joint J*. 2020;102-B(1):108-116. doi:10.1302/0301-620X.102B1.BJJ-2019-0510.R1
- Motesharei A, Rowe P, Blyth M, Jones B, Maclean A. A comparison of gait one year post operation in an RCT of robotic UKA versus traditional Oxford UKA. *Gait Posture*. 2018;62:41-45. doi:10.1016/j.gaitpost.2018.02.029
- Borus T, Roberts D, Fairchild P, Pirtle K, Baer M. Early functional performance of unicompartmental knee arthroplasty compared to total knee arthroplasty. Presented at: 2nd World Arthroplasty Congress (WAC); April 19-21, 2018. Rome, Italy.
- van der List JP, Chawla H, Villa JC, Pearle AD. Different optimal alignment but equivalent functional outcomes in medial and lateral unicompartmental knee arthroplasty. *Knee*. 2016;23(6):987-995. doi:10.1016/j.knee.2016.08.008
- Augar MA, Plate JF, Bracey DN, Jinnah A, Poehling GG, Jinnah RH. Robotic lateral and medial unicompartmental knee arthroplasty. *Oper Tech Orthop*. 2015;25(2):95-103. doi:10.1053/j.oto.2015.02.002
- Zuiderbaan HA, Khamaisy S, Thein R, Nawabi DH, Pearle AD. Congruence and joint space width alterations of the medial compartment following lateral unicompartmental knee arthroplasty. *Bone Joint J*. 2015;97-B(1):50-55. doi:10.1302/0301-620X.97B1.33057
- Thein R, Zuiderbaan HA, Khamaisy S, Nawabi DH, Poultsides LA, Pearle AD. medial unicompartmental knee arthroplasty improves patellofemoral congruence: a possible mechanistic explanation for poor association between patellofemoral degeneration and clinical outcome. *J Arthroplasty*. 2015;30(11):1917-1922. doi:10.1016/j.arth.2015.05.034
- Khamaisy S, Zuiderbaan HA, van der List JP, Nam D, Pearle AD. Medial unicompartmental knee arthroplasty improves congruence and restores joint space width of the lateral compartment. *Knee*. 2016;23(3):501-505. doi:10.1016/j.knee.2016.02.012
- Burger JA, Dooley MS, Kleebad LJ, Zuiderbaan HA, Pearle AD. What is the impact of patellofemoral joint degeneration and malalignment on patient-reported outcomes after lateral unicompartmental knee arthroplasty? [published correction appears in *Bone Joint J*. 2020 Aug;102-B(8):1108]. *Bone Joint J*. 2020;102-B(6):727-735. doi:10.1302/0301-620X.102B6.BJJ-2019-1429.R1
- Farr J 2nd, Barrett D. Optimizing patellofemoral arthroplasty. *Knee*. 2008;15(5):339-347. doi:10.1016/j.knee.2008.05.008
- Cannon A, Stolley M, Wolf B, Amendola A. Patellofemoral resurfacing arthroplasty: literature review and description of a novel technique. *Iowa Orthop J*. 2008;28:42-48.
- Odgaard A, Madsen F, Kristensen PW, Kappel A, Fabrin J. A randomized clinical trial on patellofemoral vs. total knee replacement for patellofemoral osteoarthritis. *Knee Society 2017 Mark Coventry, MD Award*. 2017 Specialty Day of the Knee Society. San Diego, CA. March 18, 2017.
- Noyes F, Barber-Westin S, Fleckenstein C, Riccobene J. Patellofemoral arthroplasty in younger patients: are recreational activities feasible? Presented at: American Academy of Orthopaedic Surgeons (AAOS) Annual Meeting; March 6-10, 2018; New Orleans, LA
- Brown NM, Sheth NP, Davis K, et al. Total knee arthroplasty has higher postoperative morbidity than unicompartmental knee arthroplasty: a multicenter analysis. *J Arthroplasty*. 2012;27(8 Suppl):86-90. doi:10.1016/j.arth.2012.03.022
- Ghomrawi HM, Eggman AA, Pearle AD. Effect of age on cost-effectiveness of unicompartmental knee arthroplasty compared with total knee arthroplasty in the U.S. *J Bone Joint Surg Am*. 2015;97(5):396-402. doi:10.2106/JBJS.N.00169

45. Slover J, Espehaug B, Havelin LI, et al. Cost-effectiveness of unicompartmental and total knee arthroplasty in elderly low-demand patients. A Markov decision analysis. *J Bone Joint Surg Am.* 2006;88(11):2348-2355. doi:10.2106/JBJS.E.01033
46. Zambianchi F, Franceschi G, Rivi E, et al. Clinical results and short-term survivorship of robotic-arm-assisted medial and lateral unicompartmental knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc.* 2020;28(5):1551-1559. doi:10.1007/s00167-019-05566-4
47. Crawford R, Varughese I, Herron E, Jaiprakash A, Donnelly W, Whitehouse S. The cost effectiveness of unicompartmental knee arthroplasty vs total knee arthroplasty. Presented at: 79th Annual Scientific Meeting of the Australian Orthopaedic Association (AOA); October 6-10, 2019; Canberra, Australia.
48. National Joint Replacement Registry. Hip, Knee & Shoulder Arthroplasty Annual Report 2018. Australian Orthopaedic Association; 2019.
49. Deese JM, Gratto-Cox G, Carter DA, Sasser TM Jr, Brown KL. Patient reported and clinical outcomes of robotic-arm assisted unicompartmental knee arthroplasty: Minimum two year follow-up. *J Orthop.* 2018;15(3):847-853. doi:10.1016/j.jor.2018.08.01850.
50. Dretakis K, Igoumenou VG. Outcomes of robotic-arm-assisted medial unicompartmental knee arthroplasty: minimum 3-year follow-up. *Eur J Orthop Surg Traumatol.* 2019;29(6):1305-1311. doi:10.1007/s00590-019-02424-4
51. Banger M. 5 year survivorship of a randomized trial of robotic arm assisted vs manual uni-compartmental knee arthroplasty. Presented at: European Knee Society (EKS) Arthroplasty Conference; May 2-3, 2019; Valencia, Spain.
52. Kazarian GS, Lonner JH, Maltenfort MG, Ghomrawi HMK, Chen AF. Cost-effectiveness of surgical and nonsurgical treatments for unicompartmental knee arthritis: a Markov model. *J Bone Joint Surg Am.* 2018;100(19):1653-1660. doi:10.2106/JBJS.17.00837
53. Clement ND, Deehan DJ, Patton JT. Robot-assisted unicompartmental knee arthroplasty for patients with isolated medial compartment osteoarthritis is cost-effective: a Markov decision analysis [published correction appears in *Bone Joint J.* 2019 Nov;101-B(11):1464]. *Bone Joint J.* 2019;101-B(9):1063-1070. doi:10.1302/0301-620X.101B9.BJJ-2018-1658.R1
54. Cool CL, Needham KA, Khlopas A, Mont MA. Revision Analysis of Robotic Arm-Assisted and Manual Unicompartmental Knee Arthroplasty. *J Arthroplasty.* 2019;34(5):926-931. doi:10.1016/j.arth.2019.01.018
55. St Mart JP, de Steiger RN, Cuthbert A, Donnelly W. The three-year survivorship of robotically assisted versus non-robotically assisted unicompartmental knee arthroplasty. *Bone Joint J.* 2020;102-B(3):319-328. doi:10.1302/0301-620X.102B3.BJJ-2019-0713.R1

**stryker**

325 Corporate Drive  
Mahwah, NJ 07430  
t: 201 831 5000

**stryker.com**

A surgeon must always rely on his or her own professional clinical judgment when deciding whether to use a particular product when treating a particular patient. Stryker does not dispense medical advice and recommends that surgeons be trained in the use of any particular product before using it in surgery.

The information presented is intended to demonstrate the breadth of Stryker's product offerings. A surgeon must always refer to the package insert, product label and/or instructions for use before using any of Stryker's products. The products depicted are CE marked according to the Medical Device Regulation 2017/745 or the Medical Device Directive 93/42/EEC. Products may not be available in all markets because product availability is subject to the regulatory and/or medical practices in individual markets. Please contact your sales representative if you have questions about the availability of products in your area.

Stryker Corporation or its divisions or other corporate affiliated entities own, use or have applied for the following trademarks or service marks: Mako, Stryker. All other trademarks are trademarks of their respective owners or holders.

MAKPKA-CG-1\_Rev-2\_25836

Copyright © 2020 Stryker.