















## References

1. ProLayer Acellular Dermal Matrix Therapy Instructions For Use.
2. Stevens PS, Stilwell R. The Biomechanics of AlloMend Acellular Dermal Matrix: Ultimate Tensile Strength. AlloSource whitepaper, M8S0062.00-01, 2014.
3. ProLayer Xenograft instructions for use, MR-4863, C
4. Data on file
5. AlloWrap Instructions for Use.
6. This information is provided for information regarding Stryker's soft tissue products. Stryker does not assume any responsibility for coding decisions, nor does it recommend codes for specific patients or procedures.
7. Delaney, R, Barrett, C, Stevens, P. AlloSource, Centennial, CO. ProChondrix Cartilage Restoration Matrix Contains Growth Factors Necessary for Hyaline Cartilage Regeneration.
8. ProChondrix CR Package Insert M1S2662.01
9. Li ST. Peripheral Nerve Repair with Collagen Conduits. *Clinical Materials* 9 (1992) 195-200.
10. In vitro data on file with Collagen Matrix, Inc.
11. Taras, John, Vipul Nanavati, and Pamela Steelman. "Nerve Conduits." *Journal of Hand Therapy* 18.2 (2005): 191-97.
12. Weber, Robert, Warren Breidenbach, Richard Brown, Michael Jabaley, and Daniel Mass. "A Randomized Prospective Study of Polyglycolic Acid Conduits for Digital Nerve Reconstruction in Humans." *Plastic and Reconstructive Surgery* 106.5 (2000): 1036-045.
13. \*The results of preclinical and in vitro studies may not be indicative of human clinical outcomes.
14. Gould, John S. Use of Collagen Conduits in Management of Painful Neuromas of the Foot. *Science, Technology, Innovation*, Dec. 2015.
15. NeuroMend Instructions for Use MS326, Rev 1.
16. Li St., Yuen D., Janssen JR., A semipermeable, Kink Resistant Type I Collagen-based Nerve Guide for PNS Repair. Collagen Matrix, Inc. 2003.
17. Animal study data on file at Collagen Matrix, Inc.
18. Neuroflex Instructions for Use MS314, Rev 2.
19. Waitayawinut T, Parisi D, Miller B, Luria S, Morton H, Chin S, Trumble T. A Comparison of Polyglycolic Acid Versus Type I Collagen Bioabsorbable Nerve Conduits in a Rat Model: An Alternative to Autografting. *Journal of Hand Surgery* 2007 Dec; Vol. 32A No. 10:1521-9.

## Trauma & Extremities

The information contained in this document is intended for healthcare professionals only. A surgeon must always rely on his or her own professional clinical judgment when deciding whether to use a particular product when treating a particular patient. Stryker does not dispense medical advice and recommends that surgeons be trained in the use of any particular product before using it in surgery.

The information presented is intended to demonstrate a Stryker product. A surgeon must always refer to the package insert, product label and/or instructions for use, including the instructions for Cleaning and Sterilization (if applicable), before using any Stryker product. Products may not be available in all markets because product availability is subject to the regulatory and/or medical practices in individual markets. Please contact your Stryker representative if you have questions about the availability of Stryker products in your area.

Stryker Corporation or its affiliates own, use, or have applied for the following trademarks or service marks: ActiShield, BioSkin, GraftJacket NOW, ProLayer, Stryker, ViaFlow. All other trademarks are trademarks of their respective owners or holders.

Content ID: AGFT-BR-6\_Rev-1, 03-2022

Copyright © 2022 Stryker

### Distributed by:

Stryker Orthopaedics  
325 Corporate Drive  
Mahwah, NJ 07430

Stryker  
1023 Cherry Road  
Memphis, TN 38117

stryker.com